

VANZEB 80WP

TECHNICAL DATA SHEET

1) Features

- Vanzeb 80WP is a contact fungicide with main active ingredient Mancozeb.
- Its formulation as a wettable powder has been developed to be used as a general use fungicide.

2) Benefits

- Can be used preventatively to control Sigatoka in Banana, Powdery Mildew and other common fungi in vegetables and other crops.
- Easy and safe to use when used in accordance with the instructions.

3) Composition

<u>Active Ingredients</u>	<u>% By Mass</u>
Mancozeb Produced from the coordination of zinc ion and manganese ethylene bis dithiocarbamate.	80 % Min (equivalent to 800g/kg)
Inert ingredients Dispersants and carrier.	20 % Max

4) Applications

- Vanzeb 80WP can be used preventatively for the control of Sigatoka in Banana and major fungal diseases of vegetables, fruit crops, ornamentals, tobacco and other field crops specified under directions for use.
- Its formulation as a wettable powder allows for it to be applied to a wide range of crops under various conditions.

5) Directions for use

GENERAL DOSAGE: Unless otherwise indicated, apply Mancozeb 80WP at the rate of 3 to 6 tbsps. (30 to 60 grams) per 16 liters water.

VANZEB 80WP

For banana follow the Plantation's standard disease control program.

<u>CROPS/DISEASES</u>	<u>APPLICATION</u>
MANGO Anthracnose, Scab	Spray after flowering. Repeat at 15 days interval until fruit development.
EGGPLANT Late blight, Fruit rot Anthracnose	When fruits appear, start spraying. Repeat at 7-10 days interval.
GRAPES Downy Mildew Black rot, Rust	Spray before and after blooms. Continue spraying through the season. Stop spraying 66 days before harvest. Treat seedlings from emergence to transplanting. In the field, spray at the rate of 9 tbsp. per 16 liters water at 7-10 days interval.
TOMATO Early and late blight Anthracnose, Leaf Spot	Start spraying when disease appears, at the rate of 5-7 tbsp. per 16 liters water. Repeat at 7-10 days interval.
POTATO Early and late blight	Start spraying when disease appears or at the 4 to 6-leaf stage. Repeat at 7-10 days interval.
ONION & GARLIC Downy Mildew Purple blotch	Spray when disease appears. Repeat as needed. Spray at 10-12 days interval or as needed.
BANANA Sigatoka, Crown rot	



INTERNATIONAL LTD

VANZEB 80WP

<u>CROPS/DISEASES</u>	<u>APPLICATION</u>
CUCUMBER Downy Mildew Anthracnose	Start spraying when runners begin to form or when disease appears. Repeat at 7 days interval.
CABBAGE, CAULIFLOWER Downy Mildew Alternaria Leafspot	Treat seedlings from emergence to transplanting at 5-6 days interval. In the field, spray at 7-10 days interval.
RICE Blast, leaf spots Sheath blight	Treat seedlings twice starting at 3 to 4-leaf stage. Spray at tillering and at flowering stages. Repeat application as needed.
TOBACCO Frog eye leaf spot	Treat seedlings from emergence to transplanting 2 to 3 times per week. In the field, spray immediately after transplanting. Repeat at 7-10 days interval.
PEPPER Anthracnose, Fruit rot	Spray when disease appears. Start spraying when disease appears. Repeat at 10-14 days interval. Stop spraying 14 days before harvest
PEANUT Cercospora leaf spot Rust	Start spraying when disease appears. Repeat at 7-10 days interval.
BEANS Rust, Downy Mildew	Seed treatment at 4 gms per kg of seeds using dry or slurry method. Cover seeds uniformly.
COTTON Damping off Blight	



VANZEB 80WP

Preparation and Application instructions

Mix the product in the appropriate application tank using the recommended dosage and required water volume. Can be used with manual ground equipment or high volume motorised and aerial equipment.

6) Supplier / Manufacturer Information

Supplied By:

AgroAdvanced International Philippines Inc.
3/F Vicente Madrigal Building
6793 Ayala Avenue, Makati City 1226

Warranty Statement:

It is a violation of FPA rules and regulations to use this product inconsistent with the label. No warranty of any kind expressed or implied is made concerning the use of this product. The user assumes all risks and liabilities resulting from improper handling, use or application.